



Last time we saw

M-th on




dual \longrightarrow

F-th on



Take M_g to be $\mathbb{R}^5 \times \mathcal{B}_4$

\longrightarrow to preserve SUSY, need
to be $\mathbb{C}Y_3$



Example: $K3$

$\mathcal{B}_4 = \mathbb{P}^1 \times \mathbb{R}^2$, i.e. F-theory on $\mathbb{R}^{1,7} \times K3$

$$K3 \text{ eq. : } y^2 = x^3 + f(u, v)xz^4 + g(u, v)z^6 \quad (*)$$

where $x, y, z, u, v \in \mathbb{C}/\sim$

$$\begin{aligned} (u, v, x, y, z) &\sim (\lambda u, \lambda v, \lambda^4 x, \lambda^6 y, z) \\ &\sim (u, v, \mu^2 x, \mu^3 y, \mu z) \end{aligned}$$

where $\mu, \lambda \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$, and $(u, v) \neq (0, 0)$,

$$(x, y, z) \neq (0, 0, 0)$$

$$\left. \begin{aligned} f(\lambda u, \lambda v) &= \lambda^8 f(u, v) \\ g(\lambda u, \lambda v) &= \lambda^{12} g(u, v) \end{aligned} \right\} \text{consistent with } (*)$$

rule: deg of $(*)$ = sum of weights

$$\lambda: 12 = 1 + 1 + 4 + 6 + 0$$

$$\mu: 6 = 0 + 0 + 2 + 3 + 1$$

→ total space is Calabi-Yau

projection $\pi: K3 \rightarrow \mathbb{P}^1: (x, y, z, u, v) \mapsto (u, v)$

$$\{ (u, v) \neq (0, 0) \mid (u, v) \sim (\lambda u, \lambda v) \}$$

at fixed $z=1, v=1$, we get

$$(*) \rightarrow y^2 = x^3 + f(u)x + g(u) \quad (**)$$

→ equation of elliptic curve in (x, y, z) space

$$\text{deg} = 6 = \text{sum of weights} = 2 + 3 + 1 = 6$$

Relation to $T^2 = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$

holomorphic coordinate: $z = x + \tau y$

$$\text{for } P \in T^2: z(P) = \int_0^P \Omega, \quad \Omega_1 = dz$$

For T^2 described by $(**)$, $\Omega_1 = \frac{c dx}{y}$, $c = \text{const.}$

then $\tau = \frac{\oint_B \Omega_1}{\oint_A \Omega_1}$

→ ambiguity in basis choice is $SL(2, \mathbb{Z})$

How to compute τ ?

Use j -invariant!

$$j(\tau) = \frac{4 \cdot (24f)^3}{\Delta}, \quad \Delta = 27g^2 + 4f^3$$

and $j(\tau)$ is $SL(2, \mathbb{Z})$ modular invariant

$$j(\tau) = e^{-2\pi i \tau} + 744 + O(e^{2\pi i \tau})$$

$\Delta = 0$ is "discriminant locus"

→ $\deg(\Delta) = 24 \rightarrow 24$ zero's on \mathbb{P}^1

→ denote by $u_i, i=1, \dots, 24$

We have: $j(\tau(u)) \sim \frac{1}{u-u_i}$ near u_i


$$\rightarrow \tau(u) \simeq \frac{1}{2\pi i} \ln(u-u_i)$$

for $u \rightarrow u_i$: $\tau \rightarrow i\infty$ (ratio of A- and B-cycle of T^2 vanishes)

Since $\tau = C_0 + \frac{i}{g_{\mathbb{H}^3}}$ this is "weak coupling"

limit : $g_{\mathbb{R}^3} \rightarrow 0$

monodromy:



$$u(\theta) = u_i + \varepsilon e^{2\pi i \theta}$$

$$\rightarrow T: \tau \mapsto \tau + 1$$

equivalently, $C_0 \mapsto C_0 + 1$, or $\oint_{u_i} F_i = \oint_{u_i} dC_0 = 1$

\rightarrow D7-brane $u = u_i$

altogether, we have 24 D7-branes

since \mathbb{P}^1 is compact, we expect $\sum_{i=1}^{24} \oint_{u_i} F_i = 0$

How can this be ?

solution: monodromies at different u_i
are related through $SL(2, \mathbb{Z})$
transformations: $T_i = M T M^{-1}$

$$M \in SL(2, \mathbb{Z})$$

\rightarrow (p, q) 7-brane related to $(1, 0)$ 7-brane
by "S-duality" transformation M .

vanishing of $pA + qB$ 1-cycle of T^2

$$\begin{pmatrix} p \\ q \end{pmatrix} = M \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

The weak coupling limit:

Want to find region with $\tau \sim \text{const}$

$$\Rightarrow \frac{f^3}{g^2} = \text{const.}$$

$$\text{solved by } g = p^3, \quad f = \overset{\text{const.}}{\alpha} p^2$$

with p a homogeneous polynomial of degree 4

\rightarrow go to coordinate patch $v=1$:

$$p = \prod_{i=1}^4 (u - u_i)$$

$$\rightarrow \Delta = (4\alpha^3 + 27) \prod_{i=1}^4 (u - u_i)^6, \quad j(\tau) = \frac{4(24\alpha)^3}{27 + 4\alpha^3}$$

$$\text{For } \alpha \sim -3/4^{1/3} \rightarrow j(\tau) \sim \infty$$

weak coupling everywhere on
base P^1


Only non-trivial $SL(2, \mathbb{Z})$ element with

$$M\tau = \frac{a\tau + b}{c\tau + d} = \tau \quad \text{is } M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\text{define } x = p\tilde{x}, \quad y = p^{3/2}\tilde{y}$$

$$\rightarrow \tilde{y}^2 = \tilde{x}^3 + \alpha\tilde{x} + 1$$

$$\rightarrow \Omega_1 = p^{-1/2} \frac{d\tilde{x}}{\tilde{y}}$$


 gives $\Omega_1 \rightarrow -\Omega_1$ and $\oint_A \Omega_1 = -\oint_A \Omega_1$
 (1) $\oint_B \Omega_1 = -\oint_B \Omega_1$

and thus $z \mapsto -z$

leads to double-cover construction of \mathbb{P}^1 :

$$X: \zeta^2 = p(u, v), \quad (u, v, \zeta) \simeq (\lambda u, \lambda v, \lambda^2 \zeta)$$

with $(u, v, \zeta) \neq (0, 0, 0)$

\rightarrow equation of 2nd elliptic curve

original \mathbb{P}^1 is recovered from X as quotient

$$X/\sigma, \quad \text{where } \sigma: \zeta \rightarrow -\zeta \quad (2)$$

when circling around u_i

\mathbb{Z}_2 transformations (1) and (2) give

together

$$K3 = (T^2 \times T^2) / \mathbb{Z}_2$$

Fixed loci of \mathbb{Z}_2 -involution σ , i.e. the u_i ,

are positions of O7-planes with D7-charge

$-4 \rightarrow 4$ D7-branes on top of O7-branes